

العدد السابع والعشرون - 02 / سبتمبر ( 2017 )

## SOME APPLICATIONS OF A LINEAR OPERATOR WITH NEGATIVE COEFFICIENTS

Aisha Ahmed Amer,<sup>1</sup> Nagat Muftah Alabbar,<sup>2</sup> Rabeaa Abd Allah  
Alshbear,<sup>3</sup> Abeer Mustafa Hasek<sup>4</sup>

<sup>1</sup> Mathematics Department, Faculty of Science -Al-Khomus, Al-Margib University. <sup>2</sup> Mathematics Department, Faculty of Education of Benghazi, University of Benghazi. <sup>3</sup> Mathematics Department, Faculty of Science -Al-Khomus, Al-Margib University. <sup>4</sup> Mathematics Department, Faculty of Science -Al-Khomus, Al-Margib University.



## SOME APPLICATIONS OF A LINEAR OPERATOR WITH NEGATIVE COEFFICIENTS

### Abstract:

The main object of this paper is to introduce and study the new subclasses  $TS_{\beta,\gamma}^{m,\delta}(\lambda,l,a,c)$  and  $TR_{\beta,\gamma}^{m,\delta}(\lambda,l,a,c)$  of analytic functions with negative coefficients defined by a linear operator. Coefficient bounds for functions belonging to these subclasses are determined. Further, an application involving fractional calculus we are also given.

### بعض التطبيقات لمعامل خطي للمعاملات سالبة

عائشة احمد عامر ، نجاه مفتاح العبار ، ربيعة عبدالله الشبير، عبير مصطفى

قسم الرياضيات بكلية التربية بنغازي ، جامعة بنغازي 2 قسم الرياضيات بكلية العلوم الخمس- جامعة المرقب 3 و 4

### ملخص.

الهدف الرئيسي من هذه الورقة البحثية هو تقديم ودراسة فئات فرعية جديدة لدوال تحليلية ذات معاملات سالبة التي تم تحديدها بمعامل خطي. تم حساب المعاملات التي تنتمي إلى هذه الفئات الفرعية. وعلاوة على ذلك، تم تطبيق هذا المعاملات على المعامل الكسري لحساب التفاضل والتكامل

## 1 Introduction

The theory of derivative and integral plays an important role in the theory of univalent functions. It is believed that Ruscheweyh (1975) was the first to give a generalized derivative operator in the theory of univalent function. Later, Salagean (1983) gave another generalized derivative operator. In the same paper, he introduced an integral operator. Many properties have been discussed and studied by many researchers for these two operators. For example, Al-Oboudi (2004) introduced a generalized Salagean operator, Al-Shaqsi and Darus (2009) generalized the operator given by Ruscheweyh (1975), while Darus and Al-Shaqsi (2008) studied both derivatives of Ruscheweyh and Salagean. These operators motivate us to create another type of derivative operator.

In this paper is to introduce and study the new subclasses of analytic functions with negative coefficients defined by a linear operator [1,2]

Let  $A(n)$  denote the class of all analytic functions in the open unit disc

$U = \{z \in \mathbb{C} : |z| < 1\}$ , of the form:

$$f(z) = z + \sum_{k=n+1}^{\infty} a_k z^k, (n \in \mathbb{N}). \quad (1)$$

Denote  $T(n)$  the subclass of  $A(n)$  consisting of functions of the form:

$$f(z) = z - \sum_{k=n+1}^{\infty} a_k z^k, (a_k \geq 0, n \in \mathbb{N}). \quad (2)$$

For functions  $f \in A(n)$  given by (1) and  $g \in A(n)$  given by  $g(z) = z + \sum_{n=2}^{\infty} b_n z^n$ , we define the Hadamard product (or convolution) of  $f$  and  $g$  by

$$(f * g)(z) = z + \sum_{k=n+1}^{\infty} a_k b_k z^k.$$

If  $f, g$  are analytic in  $U$ , we say that  $f$  is subordinate to  $g$ , denoted by  $f \prec g$ , if there exists a function  $w$  analytic in  $U$  with  $w(0) = 0$  and  $|w(z)| < 1$  ( $z \in U$ ), such that  $f(z) = g(w(z))$ , ( $z \in U$ ). It is known that  $f(z) \prec g(z)$  ( $z \in U$ )  $\Rightarrow f(0) = g(0)$  and  $f(U) \subset g(U)$ .

Let the function  $\varphi(a, c; z)$  be given by

$$\varphi(a, c; z) = \sum_{n=0}^{\infty} \frac{(a)_k}{(c)_k} z^{k+1}, (z \in U, c \neq 0, -1, -2, -3, \dots),$$

where  $(x)_k$  denotes the Pochhammer symbol (or the shifted factorial).

العدد السابع والعشرون - 02 / سبتمبر ( 2017 )

Corresponding to the function  $\varphi(a,c;z)$ , Carlson and Shaffer [12] introduced a linear operator  $L(a,c)$  by

$$L(a,c)f(z) := \varphi(a,c;z) * f(z) = \sum_{n=0}^{\infty} \frac{(a)_k}{(c)_k} a_k z^{k+1}.$$

Note that:

$L(a,a)$  is the identity operator,

and  $L(a,c) = L(a,b)L(b,c)$  ( $b,c \neq 0, -1, \dots$ ).

The author [1, 2] has recently introduced a new linear operator  $D_l^{m,\lambda}(a,b)f(z)$  as the following:

**Definition 1.1** Let

$$\phi_l^{m,\lambda}(a,b;z) = \sum_{k=0}^{\infty} \left( \frac{1+\lambda k+l}{1+l} \right)^m \frac{(a)_k}{(b)_k} z^{k+1},$$

where ( $z \in U, b \neq 0, -1, -2, -3, \dots$ ),  $\lambda \geq 0, m \in \mathbb{Z}, l \geq 0$ , and  $(x)_k$  is the Pochhammer symbol.

We defines a linear operator  $D_l^{m,\lambda}(a,b): A \rightarrow A$  by the following Hadamard product:

$$D_l^{m,\lambda}(a,b)f(z) := \phi_l^{m,\lambda}(a,b;z) * f(z) = \sum_{k=0}^{\infty} \left( \frac{1+\lambda k+l}{1+l} \right)^m \frac{(a)_k}{(b)_k} a_k z^{k+1}. \quad (3)$$

Note that:

$$D_0^{0,\lambda}(a,b)f(z) = L(a,b)f(z),$$

$$(1+l)D_l^{m,\lambda}(a,b)f(z) = (1-\lambda+l)L(a,b)f(z) + \lambda z (L(a,b)f(z))' = D_\lambda(L(a,b)f(z)), \lambda \geq 0,$$

$$D_l^{m,\lambda}(a,b)f(z) = D_\lambda(D_l^{m-1,\lambda}(a,b)f(z)), \text{ where } m \in \mathbb{N} = \{1, 2, 3, \dots\}.$$

Special cases of this operator includes:

- $D_0^{m,0}(a,b)f(z) = D_l^{0,\lambda}(a,b)f(z) = L(a,b)f(z).$

العدد السابع والعشرون - 02 / سبتمبر ( 2017 )

• the Ruscheweyh derivative operator [8] in the cases:  
 $D_0^{0,0}(\beta+1,1)f(z) = D^\beta f(z); \beta \geq -1$ .

• the Salagean derivative operator [10]:  $D_0^{m,1}(1,1)f(z)$ .

• the generalized Salagean derivative operator introduced by Al-Oboudi [9]:  
 $D_0^{m,\lambda}(1,1)f(z)$ .

• the Catas derivative operator [15]:  $D_l^{m,\lambda}(1,1)f(z)$ , and finally

• The fractional operator introduced by Owa and Srivastava [6]

$$D_0^{0,0}(2,2-\gamma)f(z) = \Omega^\gamma f(z) = \Gamma(2-\gamma)z^{-\gamma} D_z^\gamma f(z);$$

$D_z^\gamma f(z)$  is the fractional derivative of  $f$  of order  $\gamma$ ;  $\gamma \neq 2,3,4,\dots$ .

Now, we introduce new subclasses of analytic functions involving our operator  $D_l^{m,\lambda}(a,b)$ .

**Definition 1.2** A function  $f \in T(n)$  is said to be in the subclass  $TS_{\beta,\gamma}^{m,\delta}(\lambda,l,a,c)$ , for  $(0 < \beta \leq 1, \gamma \in \mathbb{C} \setminus \{0\})$  if and only if:

$$\left| \frac{1}{\gamma} \left( \frac{zv'}{v} - 1 \right) \right| < \beta, \quad (4)$$

where

$$\frac{zv'}{v} = \frac{z(D_l^{m,\lambda}(a,b))' + \delta z^2(D_l^{m,\lambda}(a,b)f(z))''}{(1-\delta)D_l^{m,\lambda}(a,b)f(z) + \delta z(D_l^{m,\lambda}(a,b)f(z))'}, \quad (5)$$

$(z \in \mathbb{U}, 0 \leq \delta \leq 1)$ .

**Definition 1.3** A function  $f \in T(n)$  is said to be in the subclass  $TR_{\beta,\gamma}^{m,\delta}(\lambda,l,a,c)$  if and only if

$$\left| \frac{1}{\gamma} (v' - 1) \right| < \beta, \quad (6)$$

$(z \in \mathbb{U}, 0 \leq \delta \leq 1, 0 < \beta \leq 1, \gamma \in \mathbb{C} \setminus \{0\})$ .

## العدد السابع والعشرون - 02 / سبتمبر ( 2017 )

We note that there are some known subclasses of  $TS_{\beta,\gamma}^{m,\delta}(\lambda,l,a,c)$  and  $TR_{\beta,\gamma}^{m,\delta}(\lambda,l,a,c)$ .

**Remark 1.1** (1) If  $m = 0$ , and  $a = c = 0$  then

$$TS_{\beta,\gamma}^{0,\delta}(\lambda,l,0,0) = S_n(\beta,\gamma,\delta).$$

(2) If  $m = 0$ , and  $a = c = 0$  then

$$TR_{\beta,\gamma}^{0,\delta}(\lambda,l,0,0) = R_n(\beta,\gamma,\delta).$$

The classes  $S_n(\beta,\gamma,\delta)$  and  $R_n(\beta,\gamma,\delta)$  were investigated in [3].

(3) If  $m = 0$ ,  $a = c = 0$  and  $\delta = 0$ ,  $\beta = |b|$ ,  $\gamma = 1$  then

$$TS_{|b|,1}^{0,0}(0,l,\lambda,c) = S_1^*(b),$$

where  $(b \in \mathbb{C} / \{0\})$ . The class  $S_1^*(b)$  was studied in [4].

## 2 Coefficient bounds

In this section, we obtain necessary and sufficient conditions for a function to be in the subclasses  $TS_{\beta,\gamma}^{m,\delta}(\lambda,l,a,c)$  and  $TR_{\beta,\gamma}^{m,\delta}(\lambda,l,a,c)$  respectively.

**Theorem 2.1** Let the function  $f$  be defined by (3). Then  $f$  belongs to the subclass  $TS_1^{m,\lambda}(a,b)$  if and only if

$$\sum_{k=n+1}^{\infty} [1 + \delta(k-1)](k + \beta|\gamma| - 1) \left| \left( \frac{\lambda(k-1) + 1 + l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} \right| a_k \leq \beta|\gamma|, \quad (7)$$

$(z \in \mathbb{U}, 0 \leq \delta \leq 1, 0 < \beta \leq 1, \gamma \in \mathbb{C} / \{0\})$ .

**Proof:** Suppose  $f \in TS_{\beta,\gamma}^{m,\delta}(\lambda,l,a,c)$ . By making use of (4) we easily obtain

$$\operatorname{Re} \left\{ \frac{zv'}{v} - 1 \right\} > -\beta|\gamma| \quad (z \in \mathbb{U}),$$

which, in view of (5), gives :

العدد السابع والعشرون - 02 / سبتمبر ( 2017 )

$$Re \left\{ \frac{- \sum_{k=n+1}^{\infty} [1+\delta(k-1)](k-1) \left( \frac{\lambda(k-1)+1+l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} a_k z^{k-1}}{\sum_{k=n+1}^{\infty} [1+\delta(k-1)] \left( \frac{\lambda(k-1)+1+l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} a_k z^{k-1}} \right\} > -\beta |\gamma|. \quad (8)$$

Setting  $z = r$  ( $0 \leq r < 1$ ) in ( 8 ) we observe that the expression in the denominator on the left hand side of ( 8 ) is positive for  $r = 0$  and also for all  $r \in (0,1)$ . Thus by letting  $r \rightarrow 1^-$ , through real values ( 8 ) leads us to the desired condition ( 7 ) of the theorem.

Conversely, by applying the hypothesis ( 8 ) and setting  $|z| = 1$ , we find by using (7)that

$$\begin{aligned} \left| \frac{z v'}{v} - 1 \right| &= \left| \frac{- \sum_{k=n+1}^{\infty} [1+\delta(k-1)](k-1) \left( \frac{\lambda(k-1)+1+l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} a_k z^k}{z - \sum_{k=n+1}^{\infty} [1+\delta(k-1)] \left( \frac{\lambda(k-1)+1+l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} a_k z^k} \right| \\ &\leq \frac{\sum_{k=n+1}^{\infty} [1+\delta(k-1)](k-1) \left( \frac{\lambda(k-1)+1+l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} a_k z^{k-1}}{1 - \sum_{k=n+1}^{\infty} [1+\delta(k-1)] \left( \frac{\lambda(k-1)+1+l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} a_k z^{k-1}} \\ &\leq \frac{\beta |\gamma| \left( 1 - \sum_{k=n+1}^{\infty} [1+\delta(k-1)] \left( \frac{\lambda(k-1)+1+l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} a_k \right)}{1 - \sum_{k=n+1}^{\infty} [1+\delta(k-1)] \left( \frac{\lambda(k-1)+1+l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} a_k} \\ &= \beta |\gamma|. \end{aligned}$$

Hence, by the maximum modulus principle, we have  $f \in TS_{\beta, \gamma}^{m, \delta}(\lambda, l, a, c)$ .

**Corollary 2.1**

Let the function  $f$  be defined by (3) and  $f \in TR_{\beta, \gamma}^{m, \delta}(\lambda, l, a, c)$ , then

$$a_k \leq \frac{\beta |\gamma|}{[1+\delta(k-1)](k+\beta|\gamma|-1) \left( \frac{\lambda(k-1)+1+l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}}}, \quad (k \geq n+1), \quad (9)$$

with equality only for functions of the form

العدد السابع والعشرون - 02 / سبتمبر ( 2017 )

$$f_z = z - \frac{\beta |\gamma|}{[1 + \delta(k-1)](k + \beta |\gamma| - 1) \left[ \left( \frac{\lambda(k-1) + 1 + l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} \right]} z^k, \quad (k \geq n+1).$$

By using the same arguments as in the proof of Theorem 2.1 we can establish the next theorem.

**Theorem 2.2** Let the function  $f$  be defined by (3). Then  $f$  belongs to the subclass  $TR_{\beta, \gamma}^{m, \delta}(\lambda, l, a, c)$  if and only if

$$\sum_{k=n+1}^{\infty} k [1 + \delta(k-1)] \left[ \left( \frac{\lambda(k-1) + 1 + l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} \right] a_k \leq \beta |\gamma|, \quad (10)$$

( $z \in U$ ,  $0 \leq \delta \leq 1$ ,  $0 < \beta \leq 1$ ,  $\gamma \in \mathbb{C} \setminus \{0\}$ ).

**Corollary 2.2**

Let function  $f$  be defined by (3) and  $f \in TR_{\beta, \gamma}^{m, \delta}(\lambda, l, a, c)$ . Then

$$a_k \leq \frac{\beta |\gamma|}{k [1 + \delta(k-1)] \left[ \left( \frac{\lambda(k-1) + 1 + l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} \right]} \quad (k \geq n+1), \quad (11)$$

with equality only for functions of the form

$$f_z = z - \frac{\beta |\gamma|}{[1 + \delta(k-1)] \left[ \left( \frac{\lambda(k-1) + 1 + l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} \right]} z^k, \quad (k \geq n+1).$$

**3 An Application of Fractional Calculus**

From among various definitions of fractional calculus (that is, fractional derivative and fractional integral), we recall here the following definitions which have been used by many authors including, for example (Owa [13], Srivastava and Owa, [14]).

**Definition 3.1** The fractional integral of order  $\eta$  is defined by

$$D_z^{-\eta} f(z) = \frac{1}{\Gamma(\eta)} \int_0^z \frac{f(t)}{(z-t)^{1-\eta}} dt,$$

where  $\eta > 0$   $f$  is an analytic function in a simply connected domain of the  $z$ -plane containing the origin and the multiplicity of  $(z-t)^{\eta-1}$  is removed by requiring  $\log(z-t)$  to be real when  $(z-t) > 0$ .

**Definition 3.2** The fractional derivative of order  $\eta$  is defined by



العدد السابع والعشرون - 02 / سبتمبر ( 2017 )

$$D_z^\eta f(z) = \frac{1}{\Gamma(1-\eta)} \frac{d}{dz} \int_0^z \frac{f(t)}{(z-t)^\eta} dt,$$

where  $0 \leq \eta < 1$   $f$  is an analytic function in a simply connected domain of the  $z$ -plane containing the origin and the multiplicity of  $(z-t)^{\eta-1}$  is removed by requiring  $\log(z-t)$  to be real when  $z-t > 0$ .

**Theorem 3.3** Let the function  $f$  defined by (3) be in the subclass  $TS_{\beta,\gamma}^{m,\delta}(\lambda, l, a, c)$ . Then for  $|z|=r < 1$ .

$$|D_z^{-\eta} f(z)| \leq$$

$$\frac{|z|^{1+\eta}}{\Gamma(2-\eta)} \left\{ 1 + \frac{\beta |\gamma| (k+1)! \Gamma(2+\eta)}{[1+\delta(k)](k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right| \Gamma(k+2+\eta)} |z|^k \right\}, \quad (12)$$

and

$$|D_z^{-\eta} f(z)| \geq$$

$$\frac{|z|^{1+\eta}}{\Gamma(2-\eta)} \left\{ 1 - \frac{\beta |\gamma| (k+1)! \Gamma(2+\eta)}{[1+\delta(k)](k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right| \Gamma(k+2+\eta)} |z|^k \right\}. \quad (13)$$

The estimates in (12) and (13) are sharps.

**Proof:**

From Definition 3.1, we get

$$\begin{aligned} \Gamma(2+\eta) z^{-\eta} D_z^{-\eta} z &= z - \sum_{k=n+1}^{\infty} \frac{\Gamma(k+1)\Gamma(2+\eta)}{\Gamma(k+1+\eta)} a_k z^k \\ &= z - \sum_{k=n+1}^{\infty} \Phi(k) a_k z^k, \end{aligned} \quad (14)$$

where

$$\Phi(k) = \frac{\Gamma(k+1)\Gamma(2+\eta)}{\Gamma(k+1+\eta)}.$$

Since  $\Phi(k)$  is a decreasing function of  $k$  we have

العدد السابع والعشرون - 02 / سبتمبر ( 2017 )

$$0 < \Phi(k) \leq \Phi(k+1) = \frac{(k+1)\Gamma(2+\eta)}{\Gamma(k+2+\eta)},$$

in view of Theorem 2.1, we have

$$\begin{aligned} & [1+\delta(k)](k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right| \sum_{k=n+1}^{\infty} a_k \leq \\ & \sum_{k=n+1}^{\infty} [1+\delta(k-1)](k+\beta|\gamma|-1) \left| \left( \frac{\lambda(k-1)+1+l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} \right| a_k \leq \beta|\gamma|, \\ & \sum_{k=n+1}^{\infty} a_k \leq \frac{\beta|\gamma|}{[1+\delta(k)](k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right|}. \end{aligned} \quad (15)$$

Using (15) and (10) we have

$$\begin{aligned} & |\Gamma(2+\eta)z^{-\eta}D_z^{-\eta}f(z)| \leq |z| + \Phi(k+1)|z|^{k+1} \sum_{k=n+1}^{\infty} a_k \leq \\ & |z| + \frac{(k+1)\Gamma(2+\eta)}{\Gamma(k+2+\eta)} \frac{\beta|\gamma|}{[1+\delta(k)](k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right|} |z|^{k+1}, \end{aligned}$$

and

$$\begin{aligned} & |\Gamma(2+\eta)z^{-\eta}D_z^{-\eta}f(z)| \geq |z| - \Phi(k+1)|z|^{k+1} \sum_{k=n+1}^{\infty} a_k \geq \\ & |z| - \frac{(k+1)\Gamma(2+\eta)}{\Gamma(k+2+\eta)} \frac{\beta|\gamma|}{[1+\delta(k)](k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right|} |z|^{k+1}, \end{aligned}$$

Which prove the inequalities of Theorem 3.3. Finally, we can easily see that the results (12) and (13) are sharp for the function  $f$  defined by

$$D_z^{-\eta}f(z) = \frac{z^{1+\eta}}{\Gamma(2-\eta)} \left\{ 1 - \frac{\beta|\gamma|(k+1)\Gamma(2+\eta)}{[1+\delta(k)](k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right| \Gamma(k+2+\eta)} z^k \right\}.$$

العدد السابع والعشرون - 02 / سبتمبر ( 2017 )

By using the same arguments as in the proof of Theorem 3.3, we can establish the next theorem.

**Theorem 3.4** Let the function  $f$  defined by (3) be in the class  $TR_{\beta,\gamma}^{m,\delta}(\lambda,l,a,c)$ . Then for  $|z| = r < 1$

$$|D_z^{-\eta}f(z)| \leq \frac{|z|^{1+\eta}}{\Gamma(2-\eta)} \left\{ 1 + \frac{\beta|\gamma|(k+1)\Gamma(2+\eta)}{(k+1)[1+\delta(k)] \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right| \Gamma(k+2+\eta)} |z|^k \right\}, \quad (16)$$

and

$$|D_z^{-\eta}f(z)| \geq \frac{|z|^{1+\eta}}{\Gamma(2-\eta)} \left\{ 1 - \frac{\beta|\gamma|(k+1)\Gamma(2+\eta)}{(k+1)[1+\delta(k)] \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right| \Gamma(k+2+\eta)} |z|^k \right\}. \quad (17)$$

The estimates in (16) and (17) are sharps.

**Theorem 3.5** Let the function  $f$  defined by (3) be in the subclass  $TS_{\beta,\gamma}^{m,\delta}(\lambda,l,a,c)$ . Then for  $|z| = r < 1$ .

$$|D_z^{\eta}f(z)| \leq \frac{|z|^{1-\eta}}{\Gamma(2-\eta)} \left\{ 1 + \frac{k\beta|\gamma|\Gamma(k+1)\Gamma(2+\eta)}{[1+\delta(k)](k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right| \Gamma(k+2+\eta)} |z|^k \right\}, \quad (18)$$

and

$$|D_z^{\eta}f(z)| \geq$$

العدد السابع والعشرون - 02 / سبتمبر ( 2017 )

$$\frac{|z|^{1-\eta}}{\Gamma(2-\eta)} \left\{ 1 - \frac{k \beta |\gamma| \Gamma(k+1) \Gamma(2+\eta)}{[1+\delta(k)](k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right| \Gamma(k+2+\eta)} |z|^k \right\}. \quad (19)$$

The estimates in (18) and (19) are sharps.

**Proof:**

From Definition 3.2, we get

$$\begin{aligned} \Gamma(2-\eta) z^\eta D_z^\eta &= z - \sum_{k=n+1}^{\infty} \frac{\Gamma(k+1) \Gamma(2-\eta)}{\Gamma(k+1+\eta)} a_k z^k \\ &= z - \sum_{k=n+1}^{\infty} k \Psi(k) a_k z^k, \end{aligned} \quad (20)$$

where

$$\Psi(k) = \frac{\Gamma(k) \Gamma(2+\eta)}{\Gamma(k+1+\eta)}.$$

Since  $\Psi(k)$  is a decreasing function of  $k$  we have

$$0 < \Psi(k) \leq \Psi(k+1) = \frac{\Gamma(k+1) \Gamma(2+\eta)}{\Gamma(k+2+\eta)},$$

in view of Theorem 2.1, we have

$$\begin{aligned} \left( \frac{1}{k} \right) [1+\delta(k)] (k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right| \sum_{k=n+1}^{\infty} k a_k &\leq \\ \sum_{k=n+1}^{\infty} [1+\delta(k-1)] (k+\beta|\gamma|-1) \left| \left( \frac{\lambda(k-1)+1+l}{1+l} \right)^m \frac{(a)_{k-1}}{(c)_{k-1}} \right| a_k &\leq \beta |\gamma|, \\ \sum_{k=n+1}^{\infty} k a_k &\leq \frac{k \beta |\gamma|}{[1+\delta(k)] (k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right|}. \end{aligned} \quad (21)$$

العدد السابع والعشرون - 02 / سبتمبر ( 2017 )

Using(21) we have

$$|\Gamma(2+\eta)z^{\eta}D_z^{\eta}f(z)| \leq |z| + \Psi(k+1)|z|^{k+1} \sum_{k=n+1}^{\infty} ka_k \leq$$

$$|z| + \frac{k\Gamma(k+1)\Gamma(2+\eta)}{\Gamma(k+2+\eta)} \frac{\beta|\gamma|}{[1+\delta(k)](k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right|} |z|^{k+1},$$

and

$$|\Gamma(2+\eta)z^{\eta}D_z^{\eta}f(z)| \geq |z| - \Psi(k+1)|z|^{k+1} \sum_{k=n+1}^{\infty} ka_k \geq$$

$$|z| - \frac{k\Gamma(k+1)\Gamma(2+\eta)}{\Gamma(k+2+\eta)} \frac{\beta|\gamma|}{[1+\delta(k)](k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right|} |z|^{k+1},$$

which prove the inequalities of Theorem 3.5. Finally, we can easily see that the results (18)and (19) are sharp for the function  $f$  defined by

$$D_z^{\eta}f(z) = \frac{z^{1-\eta}}{\Gamma(2-\eta)} \left\{ 1 - \frac{k\beta|\gamma|\Gamma(k+1)\Gamma(2+\eta)}{[1+\delta(k)](k+\beta|\gamma|) \left| \left( \frac{\lambda(k)+1+l}{1+l} \right)^m \frac{(a)_k}{(c)_k} \right| \Gamma(k+2+\eta)} z^k \right\}.$$

Many other work on analytic functions related to derivative operator and integral operator can be read in [16] , [17] and [18] .

## References

- [1] Aisha Ahmed Amer , Second Hankel Determinant for New Subclass Defined by a Linear Operator, Springer International Publishing Switzerland 2016, Chapter 6 .
- [2] Aisha Ahmed Amer and Maslina Darus, A distortion theorem for a certain class of Bazilevic function , *Int. Journal of Math. Analysis*, 6 (2012),12, 591-597.
- [3] O.Altintas, O.Ozkan, H. M. Srivastava, Neighborhoods of a class of analytic functions with negative coefficients, *Applied Math. Letters*. 13(2000), 63-67, .
- [4] A. Y. Lashin, Starlike and convex functions of complex order involving a certain linear operator, *Indian J. Pure and Appl.* 34(7) (2003), 1101-1108, .

العدد السابع والعشرون - 02 / سبتمبر ( 2017 )

- [5] J. E. Littlewood, On inequalities in the theory of functions, *Proc. London Math. Soc.* 23 481-519 (1925).
- [6] S. Owa and H. M. Srivastava, Some applications of the generalized Libera operator, *Proc. Japan Acad. Set. A Math. Sci.* 62 (1986), 125-128, .
- [7] T. M. Flett, The dual of an inequality of Hardy and Littlewood and some related inequalities, *J. Math. Anal. Appl.* 38 (1972), 746-765.
- [8] St. Ruscheweyh , New criteria for univalent functions, *Proc. Amer. Math.Soc.* 49, (1975), 109-115, .
- [9] F.M. Al-Oboudi, On univalent functions defined by a generalized Salagean Operator, *Int. J. Math. Math. Sci.* 27 (2004), 1429-1436 .
- [10] G. S. Salagean , Subclasses of univalent functions, *Lecture Notes in Math Springer-Verlag.* 1013 (1983), 362-372 .
- [11] M. Darus and K. Al-Shaqsi , Differential sandwich theorem with generalized derivative operator, *Int. J. Math. Sci.* 2(2) (2008), 75-78.
- [12] B.C. Carlson, D.B. Shaffer, Starlike and prestarlike hypergeometric functions , *SIAM J. Math. Anal.* 15(4) (1984), 737-745 .
- [13] H. M. Srivastava and S. Owa, An application of the fractional derivative, *Mathematica Japonica.* 29(3), (1984), 383-389 .
- [14] S. Owa, On the distortion theorems, *Kyungpook Math. J*,18, (1978),53-59 .
- [15] Catas.A, On a Certain Differential Sandwich Theorem Associated with a New Generalized Derivative Operator. *General Mathematics*, 4,2009,83-95.
- [16] Aisha Ahmed Amer, Rabeaa Abd Allah Alshbear & Nagat Muftah Alabbar ,Some Applications Of A Linear Operator To A Certain Subclasses Of Analytic Functions With Negative Coefficients. *Journal of Education University Al-Khomus, Al-Margib University* 10(2017).441-454. -
- [17] Aisha Ahmed Amer and Maslina Darus, Some properties for a linear operator, *AIP Conference Proceedings* 1522, 908 (2013).
- [18] Nagat.M. Mustafa & Maslina Darus, M. 2011. On a subclass of analytic functions with negative coefficient associated to an integral operator involving Hurwitz- Lerch Zeta function, "Vasile Alecsandri" University of Bacau Faculty of Sciences Scientific Studies and Research Series Mathematics and Informatics. 21( 2) : 45 - 56.